

Review Article

Assessing the Nutritional-Value-Based Therapeutic Potentials and Non-Destructive Approaches for Mulberry Fruit Assessment: An Overview

Muhammad Faisal Manzoor ¹, Abid Hussain,² Diana Tazeddinova,^{3,4}
Aizhan Abylgazinova,^{4,5} and Bin Xu ¹

¹School of Food and Biological Engineering, Jiangsu University, Zhenjiang, Jiangsu, China

²Department of Agriculture and Food Technology, Karakoram International University, Gilgit, Pakistan

³Department of Technology and Catering Organization, South Ural State University, Chelyabinsk, Russia

⁴Higher School of Technologies of Food and Processing Productions,
Zhangir Khan West Kazakhstan Agrarian Technical University, Uralsk, Kazakhstan

⁵Scientific-Production Center of Livestock and Veterinary Medicine, Nur-Sultan, Kazakhstan

Correspondence should be addressed to Bin Xu; food_oil@126.com

Received 11 January 2022; Accepted 15 February 2022; Published 24 March 2022

Academic Editor: Deepika Koundal

Copyright © 2022 Muhammad Faisal Manzoor et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Among different fruits, mulberry is the most highlighted natural gift in its superior nutritional and bioactive composition, indispensable for continuing a healthy life. It also acts as a hepatoprotective immunostimulator and improves vision, antimicrobial, anti-cancer agent, anti-stress activity, atherosclerosis, neuroprotective functions, and anti-obesity action. The mulberry fruits also help reduce neurological disorders and mental illness. The main reason for that is the therapeutic potentials present in the nutritional components of the mulberry fruit. The available methods for assessing mulberry fruits are mainly chromatographic based, which are destructive and possess many limitations. However, recently some non-invasive techniques, including chlorophyll fluorescence, image processing, and hyperspectral imaging, were employed to detect various mulberry fruit attributes. The present review attempts to collect and explore available information regarding the nutritional and medicinal importance of mulberry fruit. Besides, non-destructive methods established for the fruit are also elaborated. This work helps encourage many more research works to dig out more hidden information about the essential nutrition of mulberry that can be helpful to resolve many mental-illness-related issues.

1. Introduction

Fruits and vegetables carry health-promoting and bioactive constituents; consequently, consumers' preference has been shifted towards their extensive consumption [1]. Chemical compositions of such food items can protect from various diseases without harming the human body. Mulberry is also a nutritious fruit cultivated 5,000 years ago in China, along with sericulture. Different health-promoting compounds such as moranoline, albafluran, albanol, morusin, kuwanol, calystegine, and hydroxymoricin that can regulate metabolic

activities efficiently have been reported [2, 3]. Explorative studies on mulberry fruit have investigated different health-promoting compounds such as moranoline, albafluran, albanol, morusin, kuwanol, calystegine, and hydroxymoricin that can regulate metabolic activities efficiently [4, 5]. In recent years, some reviews were published based on the health benefits of mulberry fruits [6, 7], and only a few studies discussed analysis methods for the mulberry fruit [8]. Moreover, with the recent global increase in demand for nutrient-dense and high-quality foods, there is a strong emphasis on the non-destructive methods of assessment

with accuracy and high sensitivity for different phytochemicals. Recently, many reviews have discussed the use of non-destructive spectral imaging and spectroscopic techniques for different food and agricultural applications such as exploring the lycopene content, ripening, and maturity of the fruits and quality assessment of alcoholic beverages and spices [9–11].

Therefore, in the current attempt, we discussed non-destructive and rapid methods for the non-invasive investigation of mulberry samples and also elaborated on the importance of mulberry fruit as a nutritional and health-promoting source.

2. Nutritional Significance of Mulberry Fruits

The occurrence of ascorbic acid, carbohydrates, proteins, fats, minerals, and vitamins (thiamine, nicotinic acid, and riboflavin) and their precursors make mulberry fruit the most nutritious agricultural product for consumers [12]. However, a broad range of topographical, climatic, and soil conditions can affect plants' nutritional and chemical status. For example, moisture can range from 80.8 ± 2.81 g/100g FW, ash can vary from 0.6 ± 0.09 g/100g DW, protein can vary from 1.46 ± 0.18 g/100g DW, and the lipid can range from 0.58 ± 0.06 g/100g DW. Likewise, crude fiber (1.2 ± 0.40 g/100g DW), carbohydrate (15.30 ± 1.27 g/100g DW), and energy (72.30 ± 3.25 kcal/100g DW) can also fluctuate [13, 14]. The main sugars are fructose ($1.7\text{--}2.11$ g/100 FW) and glucose ($1.7\text{--}2.44$ g/100 FW) in mulberry fruit. The pH and the total soluble solids (TSS) in mulberry fruit range from 3.23 to 3.42 and 6.19 to 9.32, respectively [13, 14].

Moreover, the fruit also contains essential minerals elements (both macro and micro), which aids in regulating metabolic mechanisms. Potassium (K), calcium (Ca), magnesium (Mg), and sodium (Na) are essential elements, while the iron (Fe), zinc (Zn), and nickel (Ni) are among the microminerals reported in different studies [14, 15]. Evaluation of essential minerals in different mulberry varieties from different zones confirmed K as the predominant element, ranging from $906.75 \pm 41.49\%$. Trace elements assessed were ranging in a reasonable amount in the fresh matter, but selenium (Se), arsenic (As), and chromium (Cr) are the least dominant elements [16].

In mulberry fruit, 18 different amino acids were also determined, among which 9 were the essential amino acid compulsory for our body. According to the protein ratio, mulberry fruit is close to the sound quality protein foods such as milk and fish. In mulberry essential, amino acid/total amino acid (EAA/TAA) ratio is about 42% [17]. Similarly, in mulberry fruit, the polyunsaturated fatty acid (PUFA) content is higher than monounsaturated and saturated fatty acids. Moreover, behenic acid (C22: 0) and palmitoleic acid (C16: 1) are essential fatty acids reported in mulberry fruits only. All the reported fatty acids in the fruit are displayed in Table 1.

Additionally, the mulberry fruit also contains riboflavin, thiamin, folate, niacin, and vitamins B-6, A, K, and E. Ascorbic acid is higher, ranging from 36 to 36.4 mg/100g [20]. Similarly, tocopherols were also revealed by Gómez-

Mejía et al. [23] in mulberry fruit, including α -tocopherol (0.73 mg/100 g FW), δ -tocopherol (2.2 mg/100 g FW), and γ -Tocopherol (25 mg/100 g FW). Besides, the fruit also carries organic acids [22], making it a valuable, healthy product, and plays a vital role in sensory properties by imparting sugar and sour taste. Acids are primarily used as an additive, specifically acidulates (malic, tartaric, ascorbic, and citric acids), anti-oxidants (malic, tartaric, and citric acids), or preservatives (benzoic and sorbic acids) [24]. Fruits' base acids have no adverse health effects because during the metabolism they are quickly oxidized. Other vitamins and organic acids present in the fruit are presented in Table 1.

3. Phytochemicals in Mulberry Fruit

Non-nutritive phenolic compounds protect from various dysfunctions with no side effects to the consumer's health [25]. The polyphenolic content reported in mulberry fruit includes flavonoids, fibers, β -carotene, anthocyanins, anthraquinone, glycosides, and oleic acid [2, 7] (Figure 1). Berries are a good source of polyphenolic and indicate the large family, categorized by the structural attribute as phenolic acids, flavonoids, tannins, stilbenes, and lignans, which are indispensable for life [7]. In mulberry fruits, the phenolic contents fluctuate with various cultivars. Beyond the cultivars, the maturity stages of mulberry fruits also have a notable influence on the phenolic values. With the increase in the maturity stage of mulberry fruit, phenolic content also enhances [12, 26].

3.1. Flavonoids. Flavonoids are among a large group of non-nutritive polyphenolic compounds that exhibits anti-oxidant attributes that play a pivotal role in curing oxidative stress-related problems such as atherosclerosis [27]. The variation in flavonoids contents in various breeds of mulberries is notable. Chinese mulberries have higher flavonoid contents (0.0024 mg/kg) than Korean (0.0006 mg/kg) [18]. Quercetin 3-O- β -D-(6''-O-malonyl) glucoside is the most important flavonoid for delivering anti-oxidant properties in the fruit [28]. The kaempferol 3-O-glucoside content was observed in mulberry to be 3.55–47.80 mg/kg of fresh weight in *Morus atropurpurea* cv Taiwanguosuang and Yuefenshen [29]. Besides, mrin (flavonoid) was also revealed to suppress cyclosporine in tissues. Cyclosporine is an effective immune suppressive agent that reduces nitric oxide creation by the activated macrophages [30]. Studies also demonstrated inhibition of the human cytochrome CYP3A process in a pooled human liver microsome system by regular consumption of black mulberry fruit juice [31]. Studies conducted on mice also confirmed anti-stress activity in black mulberry juice due to its valuable phytochemical composition [32].

3.2. Anthocyanins. Anthocyanins are color imparting pigments, widely distributed in agricultural products including fruits, vegetables, flowers, and others [33]. Numerous studies confirmed the presence of health-promoting anthocyanins




FIGURE 1: Main mulberry functional components and their therapeutic properties.

TABLE 1: Fatty acid composition, vitamins contents, and organic acids profile in mulberry fruit.

Fatty acid				
Name	Concentration	Name	Concentration	References
Myristic acid	0.47–0.49	Behenic acid	1.3–1.37	[18, 19]
Palmitoleic acid	0.38–0.40	Tetracosanoic acid	1.0–1.04	
Palmitic acid	20–22.26	9-Octadecynoic acid	—	
Heptadecanoic acid	0.26–0.28	10-Nonadecenoic acid	—	
Linoleic acid	26–26.45	Azeloic acid	0.23	
Oleic acid	10.00–10.68	Oxiraneoctanoic acid	0.41	
Stearic acid	8–8.62	11-Eicosenoic acid	—	
Eicosanoic acid	2.10–2.45	Brassicidic acid	0.83	
Linolenic acid	0.66	PUFA	74.11–79.52	
MUFA	5.92–6.89	SFA	14.56–19.82	
Vitamin				
Name	Concentration	Name	Concentration	References
Thiamin (B ₁)	0.026–0.029 mg/100 g	Folate DFE ^b	6.00 µg/100 g	[14, 20]
Riboflavin (B ₂)	0.900–0.101 mg/100 g	Vitamin A, RAE ^c	1.00 µg/100 g	
Nicotinic acid	0.700–0.800 mg/100 g	Vitamin A, IU ^a	25 IU/100 g	
Ascorbic acid	36.00–36.40 mg/100 g	Vitamin E (α-tocopherol)	0.80–0.87 mg/100 g	
Niacin	0.600–0.620 mg/100 g	Vitamin K	7.60–7.80 µg/100 g	
Vitamin B-6	0.030–0.050 mg/100 g			
Organic acids				
Name	Concentration	Name	Concentration	References
Malic acid	9.095 mg/g	Citric acid	1.805 mg/g	[21, 22]
Tartaric acid	0.145 mg/g	Oxalic acid	0.660 mg/g	
Fumaric acid	0.213 mg/g	Succinic acid	2.836 mg/g	
Lactic acid	0.662 mg/g	Acetic acid	0.053 mg/g	

*Fatty acids results: Results are described as percentage over the total peak area content of gas chromatography-mass spectrometry analysis.

^aIU = international unit, ^bDFE; dietary folate equivalents, and ^cRAE; retinol activity equivalents.

in mulberry fruit juice. Among them, cyanidin-3-O-glucoside and cyanidin-3-rutinoside are widely distributed (Table 2) [41].

Moreover, cyanidin 3-O-β-D-glucopyranoside isolated from mulberry fruits inhibited the cerebral ischemic damage caused by oxygen-glucose deprivation in PC12 cells [42]. Similarly, anthocyanins from the mulberry fruit (black) were

also reported to prevent Cu-induced peroxidation of liposomes [42]. Meanwhile, the co-oxidation of linoleic acid and β-carotene confirmed that the extracts of mulberry (*Morus nigra*) fruits show safeguard action against peroxidative damage to biomolecules and their membranes [43]. Furthermore, Jiang et al. [44] and Wu et al. [45] concluded that the fruit anthocyanin and water extract could scavenge free

TABLE 2: Composition of flavonoids in mulberry fruit.

Class	Subclass	Compound	Contents	References
Flavonoids	Anthocyanins	Cyanidin 3-O-(6"-O- α -rhamnopyranosyl- β -D-glucopyranoside)	57 mg/g CMA	[34]
		Cyanidin-3-rutinoside	108.78 mg/g MAE	[35]
		Cyanidin-3-glucoside	301.74 mg/g MAE	[36]
		Cyanidin 3-O-(6"-O- α -rhamnopyranosyl- β -D-glucopyranoside)	270 mg/g CMA	[34]
		Cyanidin 3-O- β -D-glucopyranoside	233 mg/g CMA	[34]
		Cyanidin 7-O- β -D-glucopyranoside	33 mg/g CMA	[34]
		Pelargonidin-3- glucoside	NA	[37]
		Pelargonidin-3-rutinoside	NA	[37]
		Petunidin 3-O- β -glucopyranoside	5.1 mg/g CEE	[38]
		Rutin	0.065–7.728 mg/100 g FW	[39]
	Flavonols	Myricetin	0.66–1.18 mg/100 g DW	[40]
		Quercetin	31.88–58.42 mg/100 g DW	[40]
		Quercetin 3-O-glucoside	1.069 mg/100 g FW	[29]
		Quercetin 3-O-rutinoside	2.869 mg/100 g FW	[29]
		Quercetin 3-O-galactoside	0.002 mg/100 g FW	[29]
		Kaempferol	0.24–1.61 mg/100 g DW	[40]
		Kaempferol 3-O-rutinoside	2.00–14.00 mg/100 g DW	[18]
		Kaempferol 3-O-glucoside	1.623 mg/100 g FW	[29]
		Catechin	309.26–750.01 mg/100 g DW	[40]
		Epicatechin	8.47–17.12 mg/100 g DW	[40]
	Flavanols	Epigallocatechin gallate	0.033–0.086 mg/100 g DW	[39]
		Procyanidin B1	59.64–224.41 mg/100 g DW	[40]
		Procyanidin B2	1.02–5.66 mg/100 g DW	[40]

MAE, mulberry anthocyanin extract; NA, not available; CEE, crude ethanol extract; CMA, crude mulberry anthocyanin; DW, dry weight; and FW, frozen weight.

radicals, prevent low-density lipoprotein oxidation, reduce blood lipid, and also be found to prevent atherosclerosis.

3.3. Flavonols and Flavanols. Flavonols and flavanols are flavonoids subgroups, and the structures of these flavonoids are almost the same but different in some positions, such as C-2, C-3, and C-4. In flavonols, a double bond exists between C-2 and C-3 and, in the C ring, the carbonyl group at C-4 compared with flavanols. Mulberry fruit consists of many flavonols such as quercetin, rutin, kaempferol, and myricetin, and the derivatives of kaempferol and quercetin are the main components. Some kaempferol in glycosylated form has been found in some cultivars of mulberry fruit, such as kaempferol 3-O-rutinoside and kaempferol 3-O-glucoside [46, 47]. Usually, the flavanols do not exist as a glycoside naturally. But, in mulberry fruit, epigallocatechin gallate, catechin, procyanidin B1, epicatechin, and procyanidin B2 have been reported (Table 2).

3.4. Phenolic Acids. Mulberry fruit consists of different types of phenolic acids. Benzoic acid and hydroxycinnamic acids are the leading derivatives that represent the phenolic acids in the fruit. Ferulic acid, cinnamic acid, chlorogenic acid, o-coumaric acid, caffeic acid, and p-coumaric acid are the leading derivatives of hydroxycinnamic acid found in the samples. The gallic acid, protocatechuic acid, hydroxybenzoic acid, and vanillic acid are the essential derivatives of benzoic acid in mulberry fruit (Table 3). In mulberry fruit, the most abundant phenolic acid is chlorogenic acid ranging from 5.3 to 17.3 mg/100 g DW [49]. Furthermore, Butkhup et al. [40] reported that cinnamic acid varied from 11.63 to

15.04 mg/100 g DW and gallic acid fluctuated from 7.34 to 23.35 mg/100 g DW, which is the most prominent phenolic acids in different cultivars of the fruit.

3.5. Polysaccharides. Polysaccharides play significant parts in pathological and physiological activities [50–52]. Different polysaccharides were purified from the mulberry fruit with hypoglycemic and anti-oxidant activities using numerous purification methods as presented in Table 4. A glycoprotein extracted from the lyophilized powder and fruit juice of mulberry fruit exhibit good anti-inflammatory and anti-apoptosis agents in rats' primary splenocytes [54].

4. Biosynthesis of Anthocyanin and Phenolic Contents in Mulberry

Anthocyanins are the phenylpropanoid metabolic pathway responsible for exhibiting red, purple, and bluish colors to the mulberry fruits. Its biosynthesis begins with amino acid phenylalanine that is converted by phenylalanine ammonia-lyase (PAL), cinnamate 4-hydroxylase (C4H), and 4-coumarate-CoA ligase (4CL) to *p*-coumaroyl-CoA (anthocyanin, flavonols, and lignins precursor). Anthocyanins (cyanidin 3-O-rutinoside) are primarily synthesized by chalcone synthase (CHS), chalcone isomerase (CHI), flavanone-3-hydroxylase (F3H), flavonoid-3'-hydroxylase (F3'H), dihydroflavonol reductase (DFR), anthocyanidin synthase (ANS), anthocyanidin 3-O-glucosyltransferase (3GT), and UDP-rhamnose: anthocyanidin-3-glucoside rhamnosyltransferase (3RT). This biosynthesis is regulated by transcription factors (TFs), including MYB and basic

TABLE 3: Composition of phenolic acid in mulberry fruit.

Class	Subclass	Compound	Content	References
Phenolic acid	Hydroxycinnamic acid	Chlorogenic acid	5.3–17.3 mg/100 g DW	[48]
		Cinnamic acid	11.63–15.04 mg/100 g DW	[40]
		p-Coumaric acid	0.024–0.142 mg/100 g DW	[39]
		o-Coumaric acid	0.015 mg/g FW	[22]
		Ferulic acid	0.057–2.949 mg/100 g DW	[39]
		Caffeic acid	1.06–8.17 mg/100 g DW	[40]
	Benzoic acid	p-Hydroxybenzoic acid	0.028–0.154 mg/100 g DW	[39]
		Protocatechuic acid	0.264–0.794 mg/100 g FW	[39]
		Gallic acid	7.34–23.35 mg/100 g DW	[40]
		Vanillic acid	0.008 mg/g FW	[22]
		Syringic acid	0.049 mg/g FW	[22]

FW, frozen weight and DW, dry weight.

TABLE 4: List of isolated polysaccharides from the mulberry fruit.

Compound name	Molecular weight	Bioactivities	References
FMAP	130	—	[53]
MP	—	Anti-apoptotic and anti-inflammatory	[54]
PMF-1	71.68	—	[55]
PMF-2	84.33	—	
PMF-3	103.17	—	
MFP	—	Hypoglycemic and anti-oxidant	[56]
MFP-1	—	Hypoglycemic and anti-oxidant	
MFP-2	—	Hypoglycemic and anti-oxidant	
MFP-1	7.9, 1.0, 0.7	Hypoglycemic and anti-oxidant	[35]
MFP-2	149, 9.3, 2.61.5	Hypoglycemic and anti-oxidant	
MFP-3	167, 5, 1.5	Hypoglycemic and anti-oxidant	
MFP-4	185, 64.4, 1.5, 0.2	Hypoglycemic and anti-oxidant	[57]
MFP3P	136.6	Hypoglycemic and anti-oxidant	
JS-MP-1	1639	Anti-obesity and immunomodulation	[58]


helix-loop-helix (bHLH) TFs and WD40-repeat proteins also [59]. Cyanidin-3-rutinoside and cyanidin-3-glucoside are the major anthocyanins isolated from mulberry fruits [59].

Among flavonoids, rutin, quercetin, and kaempferol are the major existing ones in mulberry fruits. Some mulberry cultivars have been reported with glycosylated forms of quercetin and kaempferol, including quercetin 3-O-rutinoside, quercetin 3-O-glucoside, quercetin 3-O-galactoside, kaempferol 3-O-glucoside, and kaempferol 3-O-rutinoside [20]. Moreover, rutin was reportedly the most abundant phenolic acid contributing approximately 44.66% of the total phenolic acid content in eleven different mulberry fruit samples [20].

5. Mulberry Homeostasis in Human Guts

The gut response was observed in evaluating anti-oxidant studies (ABTS and FRAP) of mulberry cultivars, whereas, on stimulation of gastrointestinal digestion, white mulberry cultivars exhibited insufficient anti-oxidant activity as compared to their black counterparts. Compared with FRAP assay, white cultivars on digestion possess better anti-oxidant capacity, while the black cultivar showed better results in the non-digested form [60]. Human gut microbiota fermentation effect on the anti-oxidant and phenolic content

was observed, where the black mulberry variety showed higher anti-oxidant activity than white in ABTS scavenging activity, while, for the FRAP assay, ferric reducing capacity showed fluctuations in results collected at 0, 2, 6, 12, and 48 hours. Similarly, the phenolic acid content decreased after fermentation in white while in black initially increased to 960.42 mg gallic acid equivalent (GAE)/100 g FW 24 hours. Still, it then significantly decreased to 543.03 GAE at 48 hours. Anthocyanins were also reportedly increased at 0 h, but with a change in pH, a structural modification was observed during fermentation producing different phenolic metabolites. After in vitro digestion, mulberry anthocyanins degraded owing to the alkaline condition of intestinal digestion. Studies showed that acidic pH was considered more stable for anthocyanins' structural integrity [60, 61]. Studies indicate the potential of digested and fermented mulberry samples in suppressing the outbreak of reactive oxygen species (ROS), as highlighted in Figure 2 [62]. Anthocyanins and catechins also act as a cellular signaling messenger to regulate the anti-oxidant enzymes and activate the Keap1/Nrf2 signaling pathway that could increase the gene expression of anti-oxidant enzymes and resultantly maintain the cellular redox balance [60, 63]. Black mulberry cultivars exhibited higher bioactive compounds with potent in vitro anti-oxidant activity and cellular ROS scavenging activity among the different tested cultivars.



Due to the health-promoting nutritional composition of mulberry fruit, its applications extend to medicine, economic enhancement, industrial by-products, clinical and domestic fields [32, 42]. Numerous authors stated that mulberry (fruit, roots, and bark) has vital importance in Chinese folk medicine, reported using since 659 AD for the treatment of different ailments such as preventing diabetes, anemia, hypotension, anti-phlogistic, hepatoprotective, diuretic, hypotensive, anti-pyretic, analgesic, expectorant, and also effective against arthritis [18, 64–67]. Furthermore, the fruit and its extracts are useful against epileptic convulsions, mental problems, and hemicranias. The fruit can also prevent asthma, vitiated conditions, rheumatism, and inflammatory issues [42].

diarrhea, aphtha, flue, cough, dyspepsia, edema, fever, headache, hypertension, and rapid healing of injuries [68]. Moreover, mulberry juice is provided in febrile disorder and malaria to reduce the body temperature because the juice (mulberry) is a natural refrigerant [69]. Similarly, the products (especially syrups and recipes) prepared using mulberry fruits effectively alleviate constipation problems, insomnia, premonitory, and apoplexy dysfunctions [31].

Moreover, the fruit is also effective in treating loss of appetite problems, flatulence, controlling intestinal parasites, and, most importantly, improving the production of body fluids. Different experiments also proved its anti-hyperlipidemic properties, hypertension preventive agent, anti-hyperglycemic, and anti-allergic properties (Table 5) [7, 32, 33, 42, 69].

Chinese pharmacopeia enlists all parts of mulberry (fruit, root bark, stem, and leaves) as a critical constituent in medicinal preparations [84, 85]. Additionally, it poses an

TABLE 5: Therapeutic properties of mulberry fruit.

Therapeutic properties	Intake type	Bioactive compounds	Health effects	References
Hypolipidemic	Mulberry freeze-dried powder	Fatty acids, dietary fiber, phenolics, anthocyanins, flavonoids, and vitamins	A significant decline in serum and liver triglyceride levels, total cholesterol, serum low-density lipoprotein cholesterol, and a decrease in the atherogenic index, while the serum high-density lipoprotein cholesterol was significantly increased	[70]
	Mulberry water extract	Phenolics, anthocyanins, flavonoids, and vitamins	Significant reduction in the levels of low-density lipoprotein, cholesterol, and triglyceride	[71]
	Mulberry freeze-dried powder	Anthocyanins	Significant reduction in the low-density lipoprotein cholesterol and total cholesterol	[72]
Anti-atherosclerotic	Mulberry water extract	Phenolics, anthocyanins, flavonoids, and vitamins	Significant decrease in severe atherosclerosis in the aorta by 42–63% Significant reduction in the glycosylated serum protein and fasting blood glucose and increase anti-oxidant enzymatic activities (SOD, CAT, GSH-Px) in streptozotocin (STZ) induced diabetic mice	[71]
	Ethyl acetate-soluble extract	Phenolic compounds (25 different types)		[73]
Anti-diabetic	Mulberry fruit polysaccharides extract	Polysaccharides	Significant reduction in fasting blood glucose, oral glucose tolerance test, fasting serum insulin levels, homeostasis model of assessment-insulin resistance, glycated serum protein, and triglycerides	[74]
	Mulberry anthocyanin extract	Anthocyanin	Significant improvement in the dysfunction in diabetic mice and mitigated insulin resistance in HepG2 cells via activation of PI3K/AKT pathways	[75]
	Ramulus mori polysaccharides extract	Ramulus mori polysacchguoarides	Blood glucose-lowering and metabolism-normalizing roles and also improvement in the function of the pancreas by inhibiting the inflammatory response and attenuating the oxidative stress in pancreas tissue	[76]
Anti-obesity	Mulberry water extracts	Gallic acid, chlorogenic acid, rutin, and anthocyanins	Regulation of lipolysis and lipogenesis, which exerted the hypolipidemic and anti-obese effects	[77]
	Mulberry leaf extract and mulberry fruit extract	Cyanidin-3-glucoside, 1-deoxynojirimycin, rutin, and resveratrol	Potential anti-obesity effects through modulation of oxidative stress and obesity-induced inflammation in high fat diet-induced obesity	[78]
Anti-tumor	Mulberry anthocyanins rich extract	Anthocyanin	Targeting c-jun and p38/p53 pathways suppress tumorigenesis and cell survival but produce apoptotic death in AGS cells	[79]
	Extract	Anthocyanin	Potent protective effect on CCl4-induced liver fibrosis in rats	[80]
Hepatoprotective	Mulberry anthocyanin extract	Anthocyanin	Occurrence of the hypolipidemic effects of mulberry anthocyanin extract via inhibition of lipid biosynthesis, phosphorylation of AMPK, and stimulation of lipolysis	[81]
Neuroprotective	Mulberry extract	Cyanidin-3-O-beta-d-glucopyranoside	Neuroprotective effects on the PC12 cells exposed to hydrogen peroxide in vitro and on cerebral ischemic damage in vivo	[82]

TABLE 5: Continued.

Therapeutic properties	Intake type	Bioactive compounds	Health effects	References
Protective against cytotoxicity and oxidative stress	Mulberry juice purification and mulberry marc purification	Total flavonols, total phenolic acids, and anthocyanins contents	Potent anti-oxidant and anti-fatigue properties	[83]

anti-aging effect and imparts positive effects on blood lipid and atherosclerosis [86]. This widely grown fruit also has corrective action against bronchitis, edema, influenza, eye infections, and nosebleeds [85]. Traditionally, mulberry can also be used to treat weakness, fatigue, premature hair falling and graying, urinary problems, tinnitus, dizziness, and hypoglycemic action [42]. In contemporary medicine, mulberry is used to prepare oral syrups, add flavor, or impart color to different drugs [87].

6.1. Anti-Oxidant Potential. Natural anti-oxidants in produce continuously inactivate the reactive species (which damage cells) and keep them in minor amounts, required for normal cell functioning [88]. The in vitro free radical's assays are the most generally employed methods in estimating the anti-oxidant potential of mulberry fruits (Table 6). Generally, the anti-oxidant activity of the whole frozen fruits was 0.21–8.15%, 50.18–86.79%, 16.53–62.83%, and 0.03–38.45 μM ascorbic acid by using a metal chelating ability, DDPH activity, superoxide anion radical scavenging methods, and FRAP activity, respectively [93]. Different experiments have concluded that fruits containing anti-oxidant compounds significantly reduce specific chronic ailments [94]. Berthollide compounds, one of the secondary metabolites, were also detected in mulberry fruits. These bioactive ingredients are free radical scavengers to protect the cell from oxidation [95]. According to another study, the fruit also strengthens the oxidation safeguarding mechanism and inactivates the red-blood-cell-damaging ingredient in diabetes-induced mice [96].

Moreover, research was conducted to compare the anti-oxidant potential of different fruits. Among the tested samples, the mulberry pulp was characterized highest anti-oxidant (ferric reducing the power of 4.11 mmol/100 g wet weight) exhibiting fruit [97]. Furthermore, a spectroscopic assessment also showed the fruit juice contains efficient scavenging characteristics against superoxide, hydroxyl, and nitric acid.

6.2. Immunostimulator. The immune system (IS) is the primary regulatory system managing the body's homeostasis and plays an essential part in developing life from childbirth to death. The IS can be balanced and guarded by using different immunostimulators. Mulberry carries a more significant amount of bioactive flavonoids, particularly anthocyanins and other bioactive compounds that play an essential role in improving the consumer's immunity [98]. *Morus alba* extracts also improved cell-mediated and humoral immunity during experimental animal studies [99].

6.3. Anti-Cancer Agent. Cancer disease is one of the main reasons for the death of both humans and animals [100]. Anthocyanins extracted from the mulberry fruit exhibit inhibitory results on migration and invasion of highly metastatic A549 carcinoma cells (human lung) in a dose-dependent method [64]. The methanolic extract of *M. alba* subdued the production of tumor necrosis factor- α (TNF- α) in macrophages. It inhibited or blocked the production of nitrogen oxide, which was LPS-activated RAW2647 [101]. Mulberry fruit extracted hydroxycinnamic acid derivatives to enhance ROS production by playing as pro-oxidants and destroying the cancer cells [102]. In another study, Huang et al. [79] proposed that mulberry anthocyanins suppressed tumorigenesis and cell survival in AGS gastric cancer xenograft model cells by attacking the c-jun and p38/p53 signaling pathways. Besides, more clinical trials and evaluations verified the curative properties of anthocyanins toward cytotoxic cells, a low-cost and readily accessible source for cancer medication, and decreased cancerous cells [103].

6.4. Hepatoprotector. The liver is one of the essential organs in the human body responsible for nutrients, growth, biochemical pathways, energy supply, and several other basic mechanisms. Hepatotoxins are dangerous elements that can harm the liver [104]. Some specific bioactive compounds in mulberry fruit, such as coumarin, flavonoids, anthocyanins, and stilbenes, were described to own hepatoprotective activity [67]. Furthermore, the hydroalcoholic extract of mulberry was verified to decrease isoniazid-produced hepatotoxicity, a specific enzyme (alanine aminotransferase and aspartate aminotransferase) deficiency ailment [105].

6.5. Atherosclerosis. Atherosclerosis is the deposition of hard yellow plaques of cholesterol in arteries' inner layers, producing heart attack or coronary thrombosis. Investigations on human health proved that dietary intake of natural anti-oxidants inhibits coronary cardiovascular diseases. Oxidation of low-density lipoprotein (LDL) and cholesterol deposition are two essential factors of atherosclerosis. Though, anti-oxidants supplementation could reduce the growth of atherosclerosis ailments [106]. Valuable bioactive compounds such as quercetin and anthocyanins are proclaimed for their shielding effects as anti-oxidant nutrients. Quercetin and its conjugates are principal representatives of the flavonol group of the mulberry; flavonols have potent inhibitory results on oxidative modification of human LDL in vitro [107]. Liu et al. [108] examined the mulberry anthocyanin extract (MAE) and mulberry water extracts (MWE) for the anti-atherosclerosis effect in vitro. The MAE and

TABLE 6: Anti-oxidant potential of mulberry fruit measured through various methods.

Species	ORAC, mmol TE g ⁻¹	ABTS, mg TE 100g ⁻¹	FRAP, mg TE 100g ⁻¹	DPPH, mg TE 100g ⁻¹	Cuprac, mg TE 100g ⁻¹	References
Black mulberry	Nr	2,788.0	1,836.0	946.0	4,046.0	[31]
	Nr	0.68–1.44	0.73–1.69	Nr	Nr	[89]
	Nr	Nr	Nr	11.5–14.5	Nr	[90]
Red mulberry	Nr	0.51–0.73	0.37–0.77	Nr	Nr	[89]
	0.301–1.728	Nr	Nr	Nr	Nr	[91]
White mulberry	Nr	Nr	Nr	29.19–44.71	Nr	[92]
	Nr	Nr	Nr	10.7–12.9	Nr	[90]
Different cultivars	Nr	0.44–1.39	Nr	Nr	Nr	[22]
	Nr	0.0384–0.2073	Nr	0.0362–0.1291	Nr	[18]
Ten cultivars of red, black, and white mulberry	Nr	1.0–325.55	Nr	1.0–160.0	Nr	[29]

Nr: not reported.

MWE scavenged the DPPH radicals and repressed the electrophoretic mobility, the production of thiobarbituric acid reactive substances, and Cu²⁺ induced ApoB fragmentation in oxidation LDL ($p < 0.05$). MAE and MWE also suppress the formation of foam cells and oxidative LDL-induced macrophage death ($p < 0.05$).

6.6. Neuroprotective. According to studies, one of the principal causes of neurodegeneration is caused by free radicals [109]. In mulberry fruit, the occurrence of cyanidin and its 3-O- β -D-glucopyranoside (Cyn3-O- β -D GP) compounds protect consumers against cerebral ischemia [7]. Cyn3-O- β -D GP is a prominent neuroprotective component of mulberry fruit extract [82]. Furthermore, Cyn3-O- β -D GP has free radical scavenging and inflammation suppressing activity and protects the brain from endothelial dysfunction [110]. Similarly, mulberry fruit extract and Cyn3-O- β -D GP can prevent reactive oxygen species production and suppress neuronal disorders. Moreover, in PC12 (oxygen-glucose-deprived) cells, Cyn3-O- β -D GP improves the viability of cells and acts as a neuroprotector against cerebral ischemia.

6.7. Hypolipidemic and Anti-Obesity Action. The anti-obesity activity of mulberries was conducted both in animal and cell models by various mechanisms (Figure 2). Obesity is defined as an elegant fat collection that increases the risk of health. Obesity hurts diabetes, hypercholesterolemia, atherosclerosis, hepatic steatosis, and hyperlipidemia and reduces the number of sugar absorption that ends in body weight. Research on mulberry extract showed the inverse relationship on the melanin-concentrating hormone (MCH) receptor, which is very helpful to reduce body weight [111]. MWE consists of polyphenols such as chlorogenic acid, gallic acid, anthocyanins, and rutin. MWE reduced visceral fat, body weight induced by a high-fat diet accompanied by hypolipidemic effects by lowering cholesterol, serum triacylglycerol, the LDL/HDL ratio, and free fatty acid. Protect the liver from impairment by lowering the hepatic lipids. In a study, MWE significantly raised the receptor α and carnitine palmitoyltransferase-1 of the hepatic peroxisome, while 3-

hydroxy-3-methylglutaryl-coenzyme A reductase and fatty acid synthases enzymes were suppressed. The results indicated that the MWE regulates lipolysis and lipogenesis that eventually imparts hypolipidemic and anti-obese effects [77]. Lately, the mulberry anti-obesity mechanism was revealed. The illustrative mulberry anthocyanin could improve the mitochondrial function via the p38-AMPK-PGC1 α pathway [112]. Moreover, mulberry pelargonidin and cyanidin controlled various obese signs in male C57BL/6 mice, including a high-fat intake [113]. Peng et al. [77] described that after six weeks of feeding with polyphenol-rich extracts of mulberry, the free fatty acids and bodyweight of high-fat intake old male hamsters were decreased.

7. Non-Destructive Techniques and Food Quality Evaluation

With the rapid increase in population and awareness, good quality food is an emerging challenge globally. Therefore, researchers focus on establishing reliable approaches for authenticating agricultural products' quality parameters, including internal and external attributes. Non-destructive powerful spectroscopic techniques have been studied for applications in milk, fish, meat, fruit, vegetable, and beverages [114–118]. The spectral imaging technique is also an accessible option, which combines digital imaging and spectroscopic techniques to provide a powerful analytical device. Such imaging approaches can deliver both spatial and spectral information simultaneously, enabling the detection of the sample down to molecular levels [11, 119].

Non-destructive assessment techniques are the central part of high-quality control functions, and they assist the different established techniques as well. Non-interruptive examination leads to the surface testing of agricultural products without any interfering technique concerning the food quality and appearance. These techniques provide data on food properties such as mechanical, chemical-physical, and structural properties. The employment of a non-destructive assessment is the most suitable way for food processing [120]. Agricultural products possess anti-oxidant attributes due to bioactive compounds such as lycopene, anthocyanins, quercetin, and polyphenols, preventing

cellular oxidation. However, these functional components are highly unstable and can be destroyed by conventional methods such as high-performance liquid chromatography, gas chromatography, thin-layer chromatography, and other techniques. Therefore, to ensure the quality assessment of such nutritious compounds, non-destructive spectroscopic and imaging methods are the best available options.

7.1. Non-Destructive Technique and Mulberry Fruit Assessment

7.1.1. Chlorophyll Fluorescence (CF). CF estimation is a non-interruptive and straightforward tool, widely applied to calculate the degree of pigment changes during ripening stages of different agricultural products [121]. In general, the light absorbed by photosynthetic organisms (using chlorophyll) can undergo three different pathways, whether it can be employed to carry out the photosynthesis process result in heat production, or reemitting fluorescence (red). All the operations occur in the competition; therefore, an increase or decrease in one pathway affects the intensity of others. Moreover, estimation of the total yield of fluorescence (chlorophyll) can provide information about changes in the power of photochemistry and heat generation (Figure 3(a)) [124, 125].

Handheld equipment is cost-effective and readily available in the market. Numerous studies have been reported recently using CF to monitor the ripening stages of fruits, including tomato fruit [124], jujube (*Ziziphus jujuba* Mill.) [126], and tobacco seeds [127].

Furthermore, in another study, CF measurements and red, green, and blue (RGB) intensity were used to investigate sugars, total phenols, ABTS cation, total flavonoid, and DPPH radical scavenging properties during different ripening stages non-destructively. The fitted relationship showed a high correlation relationship between CF and RGB intensities with the tested parameters. A high correlation between CF and tested parameters was estimated from 0.82 to 0.94 during the 4–7 ripening stages, while the correlation between RGB and internal tested parameters (R^2) fluctuated from 0.93 to 0.97 for stages 4–7. The study concluded that the CF and RGB intensity values could non-destructively and rapidly assess the quality of fruits during different ripening stages [128].

7.1.2. Image Processing (IP). In recent years, a combination of machines proved promising in different research areas. For example, machine vision integrated with artificial intelligence delivered the best results for identifying and classification quality attributes of agricultural commodities [123, 129]. Besides, IP and machine vision also aid in the quality control of food items with high accuracy and rapidness and in a non-destructive manner [130, 131]. Numerous research works have been conducted using machine vision efficiently on different fruits and vegetables such as date [132], carrot [133] apples [134], banana [135], potato [136], olive [137], and pomegranate [138]. The main

components of a typical IP and essential steps are presented in Figure 3(b).

Similarly, a study was also designed for grading mulberry fruits based on maturity (ripe, unripe, and overripe) using IP and classification methods. Each segmented sample's color and textural attributes were extracted by employing a correlation-based feature selection subset (CFS) and consistency subset (CONS) as two different feature reduction models. Simultaneously, artificial neural networks (ANN) and support vector machines (SVM) were helpful in the sample classification. ANN classification combined with the CFS subset feature extraction method delivered accuracy of 100%, 100%, and 99.1% and the least mean square error (MSE) calculations of 9.2×10^{-10} , 3.0×10^{-6} , and 2.9×10^{-3} for training, validation, and test sets, respectively. Moreover, the ANN approach integrated with the CONS subset feature extraction approach delivered an acceptable model with the accuracy recorded as 100%, 98.9%, and 98.3%, and MSE resulted as 4.9×10^{-9} , 3.0×10^{-3} , and 3.1×10^{-3} for training, validation, and test sets, respectively, in a study [138].

7.1.3. Hyperspectral Imaging (HSI). A combination of spectral and imaging tools proved promising in numerous food applications recently. The method's sensitivity is due to the generation of three-dimensional data cubes by converting spectral information into spatial data. Hence, HSI can provide a spatial map composed of spectral variations at each pixel [139]. The vibrational attributes of C–H, H–O, C–O, and N–H bonds in the food system can be easily studied by employing the HSI system [140]. Compared to traditional computer vision and human vision, the HSI system has natural advantages that can highlight some of the problematic or impossible features to extract with conventional computer vision systems [141, 142]. With the advancement in optical sensing and imaging approaches, the HSI system has recently become a scientific and effective tool for monitoring and evaluating the quality of fruits and vegetables. The main components of a typical HSI are presented in Figure 3(c).

Due to its high sensitivity and non-destructive nature, HSI was employed to detect different constituents in complex food matrices. For example, visible and near-infrared (Vis-NIR) HSI (covering 400–1,700 nm range) has been investigated for non-destructive pectin polysaccharides detection in Dashi and Guihuami intact mulberry varieties. Also, four types of pectins (DASP, WSP, CSP, and TSP) were stored in the room and at low temperatures to analyze the prediction efficiency. Results revealed easier detection of pectins in Dashi samples than others, and the samples stored at room temperature delivered good prediction compared to low temperature stored samples [143]. Similarly, total anthocyanin values and anti-oxidant attribute was also determined by Huang et al. [144] in mulberry fruit using the Vis-NIR HSI method. The best prediction method for total anthocyanins and anti-oxidant property showed R^2_{val} of 0.959 and 0.995, and RPD was calculated as 4.96 and 14.25, respectively. Moreover, the non-destructive method was also used to monitor various pectins in mulberry fruits at




FIGURE 3: (a) Working mechanism of CF [122], (b) imaging system for mulberry classification [123], and (c) main component of HSI system [1].

different storage temperatures. Dilute alkali-soluble pectin (23.52–91.78 g/kg), water-soluble pectin (17.33–117.44 g/kg), chelator soluble pectin (22.91–135.52 g/kg), and total soluble pectin (63.77–344.75 g/kg) were analyzed in the study. The best prediction outcomes were recorded from dilute alkali-soluble pectin and total soluble pectins in the Dashi cultivar stored at room temperature, delivering satisfactory residual predictive deviation results of 2.31 and 1.93, respectively [145].

8. Conclusion and Future Prospects

Investigations on mulberry fruit's health-promoting bioactive compounds proved the disease-fighting attributes in treating various chronic dysfunctions. Clinical trials proved its positive impact against cardiovascular problems, HIV, diabetes, different type of cancers, and obesity and that it can prevent body cell damage, strengthen the nervous system, and can also alleviate many other chronic ailments. Despite these efforts, future work can be focused on the detection of new phytochemicals with more effective and green extraction methods, such as ultrasonication, supercritical fluid extraction, cold plasma method, low polarity water, pulsed electric field, and their integration with other non-thermal methods. In addition, due to the rich phytochemicals and anti-oxidants, the fruit is more worthy for dieticians and health care industries in future research domains. However, some of the polyphenols mechanism

in the human body is still not clear yet, need to be thoroughly studied in future works. Furthermore, the stability of polyphenols is also a challenge and further work can be addressed by proposing novel methods in enhancing their stability.

Likewise, fast, green, label-free, and non-destructive methods are also required for accurate and non-interruptive assessment of mulberry fruit attributes. Numerous imaging methods (such as soft X-ray imaging, laser backscattering imaging, multispectral imaging, resonance imaging, thermal imaging, microwave imaging, and others) and spectroscopic approaches such as surface enhance Raman spectroscopy, near-infrared spectroscopy, Fourier transforms infrared spectroscopy, and others may be the smart choice in upcoming projects.

Data Availability

The data set supporting the conclusions of this article is included within the article.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Authors' Contributions

All the authors equally contributed to this article.

Acknowledgments

This research was funded by National Natural Science Foundation of China (Grant no. 31771900).

References

- [1] A. Hussain, H. Pu, and D.-W. Sun, "Innovative nondestructive imaging techniques for ripening and maturity of fruits - a review of recent applications," *Trends in Food Science & Technology*, vol. 72, pp. 144–152, 2018.
- [2] G. D'urso, J. J. Mes, P. Montoro, R. D. Hall, and R. C. De Vos, "Identification of bioactive phytochemicals in mulberries," *Metabolites*, vol. 10, no. 1, p. 7, 2020.
- [3] E. Gómez-Mejía, C. L. Roriz, S. A. Heleno et al., "Valorisation of black mulberry and grape seeds: chemical characterization and bioactive potential," *Food Chemistry*, vol. 337, Article ID 127998, 2021.
- [4] I. Kim and J. Lee, "Variations in anthocyanin profiles and antioxidant activity of 12 genotypes of mulberry (*Morus* spp.) fruits and their changes during processing," *Antioxidants*, vol. 9, no. 3, 242 pages, 2020.
- [5] S. Shreelakshmi, M. S. Nazareth, S. S. Kumar, P. Giridhar, K. H. Prashanth, and N. P. Shetty, "Physicochemical composition and characterization of bioactive compounds of mulberry (*Morus indica* L.) fruit during ontogeny," *Plant Foods for Human Nutrition*, vol. 76, pp. 1–7, 2021.
- [6] S. Parida, K. Rayaguru, and J. Panigrahi, "Mulberry cultivation and its phytochemical benefits: a review," *Journal of Natural Remedies*, vol. 21, no. 5, pp. 33–48, 2020.
- [7] V. K. Ramappa, D. Srivastava, P. Singh et al., "Mulberries: a promising fruit for phytochemicals, nutraceuticals, and biological activities," *International Journal of Fruit Science*, vol. 20, pp. 1–26, 2020.
- [8] P. Wen, T.-G. Hu, R. J. Linhardt, S.-T. Liao, H. Wu, and Y.-X. Zou, "Mulberry: a review of bioactive compounds and advanced processing technology," *Trends in Food Science & Technology*, vol. 83, pp. 138–158, 2019.
- [9] M. Arslan, H. E. Tahir, M. Zareef et al., "Recent trends in quality control, discrimination and authentication of alcoholic beverages using nondestructive instrumental techniques," *Trends in Food Science & Technology*, vol. 107, pp. 80–113, 2021.
- [10] M. Arunkumar, A. Rajendran, S. Gunasri, M. Kowsalya, and C. Krithika, "Non-destructive fruit maturity detection methodology-A review," *Materials Today Proceedings*, 2021.
- [11] A. Hussain, H. Pu, and D.-W. Sun, "Measurements of lycopene contents in fruit: a review of recent developments in conventional and novel techniques," *Critical Reviews in Food Science and Nutrition*, vol. 59, no. 5, pp. 758–769, 2019.
- [12] S. Nayab, K. Razzaq, S. Ullah et al., "Genotypes and harvest maturity influence the nutritional fruit quality of mulberry," *Scientia Horticulturae*, vol. 266, Article ID 109311, 2020.
- [13] F. Koyuncu, M. Çetinbas, and İ. Erdal, "Nutritional constituents of wild-grown black mulberry (*Morus nigra* L.)," *Journal of Applied Botany and Food Quality*, vol. 87, 2014.
- [14] S. M. Paunović, P. Mašković, and M. Milinković, "Determination of primary metabolites, vitamins and minerals in black mulberry (*Morus nigra*) berries depending on altitude," *Vegetable Oils in Food Technology: Composition, Properties Uses*, vol. 62, no. 3, pp. 355–360, 2020.
- [15] B. Ma, Y. Luo, L. Jia et al., "Genome-wide identification and expression analyses of cytochrome P450 genes in mulberry (*Morus notabilis*)," *Journal of Integrative Plant Biology*, vol. 56, no. 9, pp. 887–901, 2014.
- [16] S. Ercisli, M. Tosun, B. Duralija, S. Voća, M. Sengul, and M. Turan, "Phytochemical content of some black (*Morus nigra* L.) and purple (*Morus rubra* L.) mulberry genotypes," *Food Technology and Biotechnology*, vol. 48, no. 1, 2010.
- [17] Y. Jiang and W.-J. Nie, "Chemical properties in fruits of mulberry species from the Xinjiang province of China," *Food Chemistry*, vol. 174, pp. 460–466, 2015.
- [18] E. M. Sánchez-Salcedo, P. Mena, C. García-Viguera, J. J. Martínez, and F. Hernández, "Phytochemical evaluation of white (*Morus alba* L.) and black (*Morus nigra* L.) mulberry fruits, a starting point for the assessment of their beneficial properties," *Journal of Functional Foods*, vol. 12, pp. 399–408, 2015.
- [19] E. M. Sánchez-Salcedo, E. Sendra, Á. A. Carbonell-Barra-china, J. J. Martínez, and F. Hernández, "Fatty acids composition of Spanish black (*Morus nigra* L.) and white (*Morus alba* L.) mulberries," *Food Chemistry*, vol. 190, pp. 566–571, 2016.
- [20] Q. Yuan and L. Zhao, "The mulberry (*Morus alba* L.) fruit-A review of characteristic components and health benefits," *Journal of Agricultural and Food Chemistry*, vol. 65, no. 48, 10383 pages, 2017.
- [21] M. R. Bozhüyök, M. Pehlivan, K. Tuncay, and B. Doğru, "Organic acid composition of selected mulberry genotypes from Aras valley," *Atatürk Üniversitesi Ziraat Fakültesi Dergisi*, vol. 46, no. 2, pp. 69–74, 2015.
- [22] M. Gundogdu, F. Muradoglu, R. I. G. Sensoy, and H. Yilmaz, "Determination of fruit chemical properties of *Morus nigra* L., *Morus alba* L. and *Morus rubra* L. by HPLC," *Scientia Horticulturae*, vol. 132, pp. 37–41, 2011.
- [23] E. Gómez-Mejía, C. L. Roriz, S. A. Heleno et al., "Valorisation of black mulberry and grape seeds: chemical characterization and bioactive potential," *Food Chemistry*, vol. 337, Article ID 127998, 2020.
- [24] S. C. Lourenço, M. Moldão-Martins, and V. D. Alves, "Antioxidants of natural plant origins: from sources to food industry applications," *Molecules*, vol. 24, no. 22, p. 4132, 2019.
- [25] M. Wang, L.-X. Gao, J. Wang et al., "Diels-Alder adducts with PTP1B inhibition from *Morus notabilis*," *Phytochemistry*, vol. 109, pp. 140–146, 2015.
- [26] T. Mahmood, F. Anwar, M. Abbas, M. C. Boyce, and N. Saari, "Compositional variation in sugars and organic acids at different maturity stages in selected small fruits from Pakistan," *International Journal of Molecular Sciences*, vol. 13, no. 2, pp. 1380–1392, 2012.
- [27] Y.-G. Liu, J.-L. Yan, Y.-Q. Ji, W.-J. Nie, and Y. Jiang, "Black mulberry ethanol extract attenuates atherosclerosis-related inflammatory factors and downregulates PPAR γ and CD36 genes in experimental atherosclerotic rats," *Food & Function*, vol. 11, no. 4, pp. 2997–3005, 2020.
- [28] S. Sarkhel, "Nutrition importance and health benefits of mulberry leaf extract: a review," *Journal of Pharmacognosy and Phytochemistry*, vol. 9, no. 5, pp. 689–695, 2020.
- [29] Q. Jin, J. Yang, L. Ma, J. Cai, and J. Li, "Comparison of polyphenol profile and inhibitory activities against oxidation and α -glucosidase in mulberry (*Genus Morus*) cultivars from China," *Journal of Food Science*, vol. 80, no. 11, Article ID C2440, 2015.
- [30] K. Thakur, Y.-Y. Zhu, J.-Y. Feng et al., "Morin as an imminent functional food ingredient: an update on its enhanced efficacy in the treatment and prevention of metabolic

- syndromes," *Food & Function*, vol. 11, no. 10, pp. 8424–8443, 2020.
- [31] M. Tomas, G. Toydemir, D. Boyacioglu, R. Hall, J. Beekwilder, and E. Capanoglu, "The effects of juice processing on black mulberry antioxidants," *Food Chemistry*, vol. 186, pp. 277–284, 2015.
 - [32] N. Kishore and D. Kumar, "Pharmacological Properties of Mulberry (*Morus Alba*)," *Assessment of Medicinal Plants for Human Health: Phytochemistry, Disease Management, Novel Applications*, vol. 37, 2020.
 - [33] W. Kalt, A. Cassidy, L. R. Howard et al., "Recent research on the health benefits of blueberries and their anthocyanins," *Advances in nutrition (Bethesda, Md.)*, vol. 11, no. 2, pp. 224–236, 2020.
 - [34] Q. Du, J. Zheng, and Y. Xu, "Composition of anthocyanins in mulberry and their antioxidant activity," *Journal of Food Composition and Analysis*, vol. 21, no. 5, pp. 390–395, 2008.
 - [35] C. Chen, L.-J. You, A. M. Abbasi, X. Fu, R. H. Liu, and C. Li, "Characterization of polysaccharide fractions in mulberry fruit and assessment of their antioxidant and hypoglycemic activities in vitro," *Food & Function*, vol. 7, no. 1, pp. 530–539, 2016.
 - [36] C. Chen, B. Zhang, X. Fu, and R. H. Liu, "A novel polysaccharide isolated from mulberry fruits (*Morus alba* L.) and its selenide derivative: structural characterization and biological activities," *Food & Function*, vol. 7, no. 6, pp. 2886–2897, 2016.
 - [37] H.-P. Huang, Y.-W. Shih, Y.-C. Chang, C.-N. Hung, and C.-J. Wang, "Chemoinhibitory effect of mulberry anthocyanins on melanoma metastasis involved in the Ras/PI3K pathway," *Journal of Agricultural and Food Chemistry*, vol. 56, no. 19, pp. 9286–9293, 2008.
 - [38] F. Sheng, Y. Wang, X. Zhao, N. Tian, H. Hu, and P. Li, "Separation and identification of anthocyanin extracted from mulberry fruit and the pigment binding properties toward human serum albumin," *Journal of Agricultural and Food Chemistry*, vol. 62, no. 28, pp. 6813–6819, 2014.
 - [39] C. Qin, Y. Li, W. Niu, Y. Ding, R. Zhang, and X. Shang, "Analysis and characterisation of anthocyanins in mulberry fruit," *Czech Journal of Food Sciences*, vol. 28, no. 2, pp. 117–126, 2010.
 - [40] L. Butkhup, W. Samappito, and S. Samappito, "Phenolic composition and antioxidant activity of white mulberry (*Morus alba* L.) fruits," *International Journal of Food Science and Technology*, vol. 48, no. 5, pp. 934–940, 2013.
 - [41] M. I. De Rosas, L. Deis, L. Martínez, M. Durán, E. Malovini, and J. B. Cavagnaro, "Anthocyanins in nutrition: biochemistry and health benefits," in *Psychiatry and Neuroscience Update*, pp. 143–152, Springer, 2019.
 - [42] A. Mandal, "Nutraceutical and medicinal property of mulberry fruits: a review on its pharmacological potential," MDPI, Basel, Switzerland, 2020.
 - [43] G. B. Escher, M. B. Marques, M. A. V. do Carmo et al., "*Clitoria ternatea* L. petal bioactive compounds display antioxidant, antihemolytic and antihypertensive effects, inhibit α -amylase and α -glucosidase activities and reduce human LDL cholesterol and DNA induced oxidation," *Food Research International*, vol. 128, Article ID 108763, 2020.
 - [44] Y. Jiang, M. Dai, W.-J. Nie, X.-R. Yang, and X.-C. Zeng, "Effects of the ethanol extract of black mulberry (*Morus nigra* L.) fruit on experimental atherosclerosis in rats," *Journal of Ethnopharmacology*, vol. 200, pp. 228–235, 2017.
 - [45] C.-S. Wu, T.-J. Chung, Y.-J. Lee, J.-D. Hsu, and H.-J. Lee, "Mulberry supplementation reduces lipid deposition and protects hamster retina from oxLDL damage," *Journal of Functional Foods*, vol. 71, Article ID 104007, 2020.
 - [46] M. L. Chávez-González, L. Sepúlveda, D. K. Verma et al., "Conventional and emerging extraction processes of flavonoids," *Processes*, vol. 8, no. 4, p. 434, 2020.
 - [47] W.-T. Ju, O.-C. Kwon, H.-B. Kim, G.-B. Sung, H.-W. Kim, and Y.-S. Kim, "Qualitative and quantitative analysis of flavonoids from 12 species of Korean mulberry leaves," *Journal of Food Science & Technology*, vol. 55, no. 5, pp. 1789–1796, 2018.
 - [48] T. Mahmood, F. Anwar, M. Abbas, and N. Saari, "Effect of maturity on phenolics (phenolic acids and flavonoids) profile of strawberry cultivars and mulberry species from Pakistan," *International Journal of Molecular Sciences*, vol. 13, no. 4, pp. 4591–4607, 2012.
 - [49] M. K. Gecer, M. Akin, M. Gundogdu, S. P. Eydurán, S. Ercisli, and E. Eydurán, "Organic acids, sugars, phenolic compounds, and some horticultural characteristics of black and white mulberry accessions from Eastern Anatolia," *Canadian Journal of Plant Science*, vol. 96, no. 1, pp. 27–33, 2016.
 - [50] I. Giavasis, "Bioactive fungal polysaccharides as potential functional ingredients in food and nutraceuticals," *Current Opinion in Biotechnology*, vol. 26, pp. 162–173, 2014.
 - [51] J.-H. Xie, M.-L. Jin, G. A. Morris et al., "Advances on bioactive polysaccharides from medicinal plants," *Critical Reviews in Food Science and Nutrition*, vol. 56, no. sup1, pp. 60–84, 2016.
 - [52] X. Xu, P. Xu, C. Ma, J. Tang, and X. Zhang, "Gut microbiota, host health, and polysaccharides," *Biotechnology Advances*, vol. 31, no. 2, pp. 318–337, 2013.
 - [53] W. Wei, W. Zhou, N. Zang, and L. Jiang, "Structural analysis of a polysaccharide from *fructus mori albae*," *Carbohydrate Polymers*, vol. 70, no. 3, pp. 341–344, 2007.
 - [54] C.-J. Liu and J.-Y. Lin, "Protective effects of strawberry and mulberry fruit polysaccharides on inflammation and apoptosis in murine primary splenocytes," *Journal of Food and Drug Analysis*, vol. 22, no. 2, pp. 210–219, 2014.
 - [55] R. Tian, "Separation, purification and composition analysis of polysaccharides from *Mori fructus*," *West China Journal of Pharmaceutical Sciences*, vol. 29, pp. 401–404, 2014.
 - [56] C. Chen, L.-J. You, A. M. Abbasi, X. Fu, and R. H. Liu, "Optimization for ultrasound extraction of polysaccharides from mulberry fruits with antioxidant and hyperglycemic activity in vitro," *Carbohydrate Polymers*, vol. 130, pp. 122–132, 2015.
 - [57] Y. Chen, W. Zhang, T. Zhao et al., "Adsorption properties of macroporous adsorbent resins for separation of anthocyanins from mulberry," *Food Chemistry*, vol. 194, pp. 712–722, 2016.
 - [58] J. S. Lee, A. Synytsya, H. B. Kim et al., "Purification, characterization and immunomodulating activity of a pectic polysaccharide isolated from Korean mulberry fruit Oddi (*Morus alba* L.)," *International Immunopharmacology*, vol. 17, no. 3, pp. 858–866, 2013.
 - [59] S. Zhao, C. H. Park, J. Yang et al., "Molecular characterization of anthocyanin and betulinic acid biosynthesis in red and white mulberry fruits using high-throughput sequencing," *Food Chemistry*, vol. 279, pp. 364–372, 2019.
 - [60] T. Bao, Y. Li, J. Xie, Z. Jia, and W. Chen, "Systematic evaluation of polyphenols composition and antioxidant activity of mulberry cultivars subjected to gastrointestinal digestion and gut microbiota fermentation," *Journal of Functional Foods*, vol. 58, pp. 338–349, 2019.

- [61] A. Castañeda-Ovando, M. d. L. Pacheco-Hernández, M. E. Pérez-Hernández, J. A. Rodríguez, and C. A. Galán-Vidal, "Chemical studies of anthocyanins: a review," *Food Chemistry*, vol. 113, no. 4, pp. 859–871, 2009.
- [62] J. Bernatoniene and D. Kopustinskiene, "The role of catechins in cellular responses to oxidative stress," *Molecules*, vol. 23, no. 4, p. 965, 2018.
- [63] Y. P. Hwang, J. H. Choi, H. J. Yun et al., "Anthocyanins from purple sweet potato attenuate dimethylnitrosamine-induced liver injury in rats by inducing Nrf2-mediated antioxidant enzymes and reducing COX-2 and iNOS expression," *Food and Chemical Toxicology*, vol. 49, no. 1, pp. 93–99, 2011.
- [64] K.-C. Cheng, C.-J. Wang, Y.-C. Chang et al., "Mulberry fruits extracts induce apoptosis and autophagy of liver cancer cell and prevent hepatocarcinogenesis in vivo," *Journal of Food and Drug Analysis*, vol. 28, no. 1, pp. 84–93, 2020.
- [65] R. A. Kadam, N. D. Dhumal, and V. B. Khyade, "The Mulberry, *Morus alba* (L.): the medicinal herbal source for human health," *International Journal of Current Microbiology and Applied Sciences*, vol. 8, no. 4, pp. 2941–2964, 2019.
- [66] I. Khalifa, D. Xia, K. Dutta, J. Peng, Y. Jia, and C. Li, "Mulberry anthocyanins exert anti-AGEs effects by selectively trapping glyoxal and structural-dependently blocking the lysyl residues of β -lactoglobulins," *Bioorganic Chemistry*, vol. 96, Article ID 103615, 2020.
- [67] C. Ren, Y. Zhang, W. Cui et al., "A polysaccharide extract of mulberry leaf ameliorates hepatic glucose metabolism and insulin signaling in rats with type 2 diabetes induced by high fat-diet and streptozotocin," *International Journal of Biological Macromolecules*, vol. 72, pp. 951–959, 2015.
- [68] H. Zhang, Z. Ma, X. Luo, and X. Li, "Effects of mulberry fruit (*Morus alba* L.) consumption on health outcomes: a mini-review," *Antioxidants*, vol. 7, no. 5, 69 pages, 2018.
- [69] M. Y. Naeem, "Medicinal potentials and health benefits of black mulberry," *Eurasian Journal of Food Science Technology*, vol. 4, no. 1, pp. 1–5, 2020.
- [70] X. Yang, L. Yang, and H. Zheng, "Hypolipidemic and antioxidant effects of mulberry (*Morus alba* L.) fruit in hyperlipidaemia rats," *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association*, vol. 48, no. 8-9, pp. 2374–2379, 2010.
- [71] C.-C. Chen, L.-K. Liu, J.-D. Hsu, H.-P. Huang, M.-Y. Yang, and C.-J. Wang, "Mulberry extract inhibits the development of atherosclerosis in cholesterol-fed rabbits," *Food Chemistry*, vol. 91, no. 4, pp. 601–607, 2005.
- [72] A. Anchalee Sirikancharon, A. Akkarach Bumrungpert, W. Wiroje Kaewruang, T. Tipanee Senawong, and P. Patcharanee Pavadhgul, "The effect of mulberry fruits consumption on lipid profiles in hypercholesterolemic subjects: a randomized controlled trial," *Journal of Pharmacy and Nutrition Sciences*, vol. 6, no. 1, pp. 7–14, 2016.
- [73] Y. Wang, L. Xiang, C. Wang, C. Tang, and X. He, "Anti-diabetic and antioxidant effects and phytochemicals of mulberry fruit (*Morus alba* L.) polyphenol enhanced extract," *PLoS One*, vol. 8, no. 7, 2013.
- [74] Y. Jiao, X. Wang, X. Jiang, F. Kong, S. Wang, and C. Yan, "Antidiabetic effects of *Morus alba* fruit polysaccharides on high-fat diet- and streptozotocin-induced type 2 diabetes in rats," *Journal of Ethnopharmacology*, vol. 199, pp. 119–127, 2017.
- [75] F. Yan, G. Dai, and X. Zheng, "Mulberry anthocyanin extract ameliorates insulin resistance by regulating PI3K/AKT pathway in HepG2 cells and db/db mice," *Journal of Nutritional Biochemistry*, vol. 36, pp. 68–80, 2016.
- [76] C. Guo, R. Li, N. Zheng, L. Xu, T. Liang, and Q. He, "Anti-diabetic effect of ramulus mori polysaccharides, isolated from *Morus alba* L., on STZ-diabetic mice through blocking inflammatory response and attenuating oxidative stress," *International Immunopharmacology*, vol. 16, no. 1, pp. 93–99, 2013.
- [77] C.-H. Peng, L.-K. Liu, C.-M. Chuang, C.-C. Chyau, C.-N. Huang, and C.-J. Wang, "Mulberry water extracts possess an anti-obesity effect and ability to inhibit hepatic lipogenesis and promote lipolysis," *Journal of Agricultural and Food Chemistry*, vol. 59, no. 6, pp. 2663–2671, 2011.
- [78] H. H. Lim, S. O. Lee, S. Y. Kim, S. J. Yang, and Y. Lim, "Anti-inflammatory and antiobesity effects of mulberry leaf and fruit extract on high fat diet-induced obesity," *Experimental Biology and Medicine*, vol. 238, no. 10, pp. 1160–1169, 2013.
- [79] H.-P. Huang, Y.-C. Chang, C.-H. Wu, C.-N. Hung, and C.-J. Wang, "Anthocyanin-rich Mulberry extract inhibit the gastric cancer cell growth in vitro and xenograft mice by inducing signals of p38/p53 and c-jun," *Food Chemistry*, vol. 129, no. 4, pp. 1703–1709, 2011.
- [80] Y. Li, Z. Yang, S. Jia, and K. Yuan, "Protective effect and mechanism of action of mulberry marc anthocyanins on carbon tetrachloride-induced liver fibrosis in rats," *Journal of Functional Foods*, vol. 24, pp. 595–601, 2016.
- [81] J.-J. Chang, M.-J. Hsu, H.-P. Huang, D.-J. Chung, Y.-C. Chang, and C.-J. Wang, "Mulberry anthocyanins inhibit oleic acid induced lipid accumulation by reduction of lipogenesis and promotion of hepatic lipid clearance," *Journal of Agricultural and Food Chemistry*, vol. 61, no. 25, pp. 6069–6076, 2013.
- [82] T. H. Kang, J. Y. Hur, H. B. Kim, J. H. Ryu, and S. Y. Kim, "Neuroprotective effects of the cyanidin-3-O- β -D-glucopyranoside isolated from mulberry fruit against cerebral ischemia," *Neuroscience Letters*, vol. 391, no. 3, pp. 122–126, 2006.
- [83] D.-Q. Jiang, Y. Guo, D.-H. Xu, Y.-S. Huang, K. Yuan, and Z.-Q. Lv, "Antioxidant and anti-fatigue effects of anthocyanins of mulberry juice purification (MJP) and mulberry marc purification (MMP) from different varieties mulberry fruit in China," *Food and Chemical Toxicology*, vol. 59, pp. 1–7, 2013.
- [84] Y. Qiu, S. Zhang, H. Yu et al., "Identification and characterization of two novel geminiviruses associated with paper mulberry (*Broussonetia papyrifera*) leaf curl disease," *Plant Disease*, 2020.
- [85] P. Sharma, "Mulberry as a life savior-a review," *Journal of Pharmacognosy and Phytochemistry*, vol. 9, no. 2, pp. 2445–2451, 2020.
- [86] J. Wattanathorn, S. Muchimapura, W. Thukhammee et al., "Mulberry fruits protects against age-related cognitive decline," *American Journal of Applied Sciences*, vol. 9, no. 9, p. 1503, 2012.
- [87] N. P. Minh and D. T. A. Dao, "Investigation of *Saccharomyces cerevisiae* in fermented mulberry juice," *International Journal of Scientific & Technology Research*, vol. 2, no. 11, pp. 329–338, 2013.
- [88] M. E. Abd El-Hack, M. T. El-Saadony, M. E. Shafi et al., "Antimicrobial and antioxidant properties of chitosan and its derivatives and their applications: a review," *International Journal of Biological Macromolecules*, 2020.
- [89] M. Özgen, S. Serçe, and C. Kaya, "Phytochemical and antioxidant properties of anthocyanin-rich *Morus nigra* and

- Morus rubra fruits," *Scientia Horticulturae*, vol. 119, no. 3, pp. 275–279, 2009.
- [90] Á. Calín-Sánchez, J. J. Martínez-Nicolás, S. Munera-Picazo, Á. A. Carbonell-Barrachina, P. Legua, and F. Hernández, "Bioactive compounds and sensory quality of black and white mulberries grown in Spain," *Plant Foods for Human Nutrition*, vol. 68, no. 4, pp. 370–377, 2013.
- [91] M. Isabelle, B. L. Lee, C. N. Ong, X. Liu, and D. Huang, "Peroxy radical scavenging capacity, polyphenolics, and lipophilic antioxidant profiles of mulberry fruits cultivated in southern China," *Journal of Agricultural and Food Chemistry*, vol. 56, no. 20, pp. 9410–9416, 2008.
- [92] W. Zhang, F. Han, J. He, and C. Duan, "HPLC-DAD-ESI-MS/MS analysis and antioxidant activities of non-anthocyanin phenolics in mulberry (morus albaL.)," *Journal of Food Science*, vol. 73, no. 6, pp. 512–518, 2008.
- [93] M. M. Natić, D. Č Dabić, A. Papetti et al., "Analysis and characterisation of phytochemicals in mulberry (Morus alba L.) fruits grown in Vojvodina, North Serbia," *Food Chemistry*, vol. 171, pp. 128–136, 2015.
- [94] A. García-Sánchez, A. G. Miranda-Díaz, and E. G. Cardona-Muñoz, "The role of oxidative stress in physiopathology and pharmacological treatment with pro-and antioxidant properties in chronic diseases," *Oxidative Medicine Cellular Longevity*, vol. 2020, 2020.
- [95] Ö. Kalkışım, "Determination of the pomological and morphological properties of white mulberry types growing in transition region between mild and continental climates," *Journal of Food Agriculture and Environment*, vol. 11, no. 1, 2013.
- [96] G. Chandrasegaran, C. Elanchezhian, K. Ghosh, and S. Sethupathy, "Berberine chloride ameliorates oxidative stress, inflammation and apoptosis in the pancreas of Streptozotocin induced diabetic rats," *Biomedicine & Pharmacotherapy*, vol. 95, pp. 175–185, 2017.
- [97] C. Guo, J. Yang, J. Wei, Y. Li, J. Xu, and Y. Jiang, "Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay," *Nutrition Research*, vol. 23, no. 12, pp. 1719–1726, 2003.
- [98] M. M. Awais and M. Akhtar, "Evaluation of some sugarcane (saccharum officinarum L.) extracts for immunostimulatory and growth promoting effects in industrial broiler chickens," *Pakistan Veterinary Journal*, vol. 32, no. 3, 2012.
- [99] S. E. Bharani, M. Asad, S. S. Dhamanigi, and G. K. Chandrakala, "Immunomodulatory activity of methanolic extract of Morus alba Linn. (mulberry) leaves," *Pakistan Journal of Pharmaceutical Sciences*, vol. 23, no. 1, pp. 63–8, 2010.
- [100] J. Ni and L. Zhang, "Cancer cachexia: definition, staging, and emerging treatments," *Cancer Management and Research*, vol. 12, pp. 5597–5605, 2020.
- [101] E. M. Choi and J. K. Hwang, "Effects of Morus alba leaf extract on the production of nitric oxide, prostaglandin E2 and cytokines in RAW264.7 macrophages," *Fitoterapia*, vol. 76, no. 7-8, pp. 608–613, 2005.
- [102] A. Trivellini, M. Lucchesini, R. Maggini et al., "Lamiaceae phenols as multifaceted compounds: bioactivity, industrial prospects and role of "positive-stress"," *Industrial Crops and Products*, vol. 83, pp. 241–254, 2016.
- [103] J. C. Jeong, S. W. Jang, T. H. Kim, C. H. Kwon, and Y. K. Kim, "Mulberry fruit (Morus fructus) extracts induce human glioma cell death in vitro through ROS-dependent mitochondrial pathway and inhibits glioma tumor growth in vivo," *Nutrition and Cancer*, vol. 62, no. 3, pp. 402–412, 2010.
- [104] D. Muhammad, N. Chand, S. Khan, A. Sultan, and M. Mushtaq, "Hepatoprotective role of milk thistle (Silybum marianum) in meat type chicken fed aflatoxin B 1 contaminated feed," *Pakistan Veterinary Journal*, vol. 32, no. 3, 2012.
- [105] F. Muhammad, M. S. Zafar, T. Khaliq, I. Javed, and M. K. Saleemi, "Nephroprotective effects of morus alba linn against isoniazid-induced toxicity in albino rabbits," *Pakistan Veterinary Journal*, vol. 34, no. 4, 2014.
- [106] S. Aydin, O. Yilmaz, and Z. Gökçe, "Protective effect of Morus nigra L.(mulberry) fruit extract on the liver fatty acid profile of Wistar rats," *Pakistan Journal of Zoology*, vol. 47, no. 1, pp. 255–261, 2015.
- [107] A. J. Day and G. Williamson, "Biomarkers for exposure to dietary flavonoids: a review of the current evidence for identification of quercetin glycosides in plasma," *British Journal of Nutrition*, vol. 86, no. 1, pp. 105–110, 2001.
- [108] L. K. Liu, H. J. Lee, Y. W. Shih, C. C. Chyau, and C. J. Wang, "Mulberry anthocyanin extracts inhibit LDL oxidation and macrophage-derived foam cell formation induced by oxidative LDL," *Journal of Food Science*, vol. 73, no. 6, 2008.
- [109] V. R. Muddapu, S. A. P. Dharshini, V. S. Chakravarthy, and M. M. Gromiha, "Neurodegenerative diseases—is metabolic deficiency the root cause?" *Frontiers in Neuroscience*, vol. 14, 2020.
- [110] C. Kaewmool, S. Udomruk, T. Phitak, P. Pothacharoen, and P. Kongtawelert, "Cyanidin-3-O-Glucoside protects PC12 cells against neuronal apoptosis mediated by LPS-stimulated BV2 microglial activation," *Neurotoxicity Research*, vol. 37, no. 1, pp. 111–125, 2020.
- [111] B. K. Oh, K.-S. Oh, K.-I. Kwon, S. Y. Ryu, Y. S. Kim, and B. H. Lee, "Melanin-concentrating hormone-1 receptor binding activity of pheophorbides isolated from morus alba leaves," *Phytotherapy Research*, vol. 24, no. 6, pp. 919–923, 2010.
- [112] Y. You, X. Yuan, H. J. Lee, W. Huang, W. Jin, and J. Zhan, "Mulberry and mulberry wine extract increase the number of mitochondria during brown adipogenesis," *Food & Function*, vol. 6, no. 2, pp. 401–408, 2015.
- [113] T. Wu, X. Qi, Y. Liu et al., "Dietary supplementation with purified mulberry (Morus australis Poir) anthocyanins suppresses body weight gain in high-fat diet fed C57BL/6 mice," *Food Chemistry*, vol. 141, no. 1, pp. 482–487, 2013.
- [114] M. L. Amodio, F. Ceglie, M. M. A. Chaudhry, F. Piazzolla, and G. Colelli, "Potential of NIR spectroscopy for predicting internal quality and discriminating among strawberry fruits from different production systems," *Postharvest Biology and Technology*, vol. 125, pp. 112–121, 2017.
- [115] Y. Lu, Y. Huang, and R. Lu, "Innovative hyperspectral imaging-based techniques for quality evaluation of fruits and vegetables: a review," *Applied Sciences*, vol. 7, no. 2, p. 189, 2017.
- [116] M. Moudache, C. Nerín, M. Colon, and F. Zaidi, "Antioxidant effect of an innovative active plastic film containing olive leaves extract on fresh pork meat and its evaluation by Raman spectroscopy," *Food Chemistry*, vol. 229, pp. 98–103, 2017.
- [117] D. Urbonaviciene and P. Viskelis, "The cis-lycopene isomers composition in supercritical CO2 extracted tomato by-products," *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, vol. 85, pp. 517–523, 2017.
- [118] K. Wang, D.-W. Sun, Q. Wei, and H. Pu, "Quantification and visualization of α -tocopherol in oil-in-water emulsion based delivery systems by Raman microspectroscopy,"

- Lebensmittel-Wissenschaft & Technologie*, vol. 96, pp. 66–74, 2018.
- [119] L.-H. Peng, S.-Y. Xu, Y.-H. Shan et al., “Sequential release of salidroside and paeonol from a nanosphere-hydrogel system inhibits ultraviolet B-induced melanogenesis in Guinea pig skin,” *International Journal of Nanomedicine*, vol. 9, p. 1897, 2014.
- [120] P. D. C. Sanchez, N. Hashim, R. Shamsudin, and M. Z. Mohd Nor, “Applications of imaging and spectroscopy techniques for non-destructive quality evaluation of potatoes and sweet potatoes: a review,” *Trends in Food Science & Technology*, vol. 96, pp. 208–221, 2020.
- [121] M. Abdelhamid, Y. A. Sudnik, and A. Y. Alipichev, “Chlorophyll fluorescence as an indicator OF fruit ripening,” *Paper Presented at the Proceedings of the International Scientific Conference of Young Scientists and Specialists Dedicated to the 150th Anniversary of AV Leontovich*, 2019.
- [122] A. N. Misra, M. Misra, and R. Singh, “Chlorophyll fluorescence in plant biology,” *Biophysics*, vol. 7, pp. 171–192, 2012.
- [123] H. Azarmdel, S. S. Mohtasebi, A. Jafari, and A. Rosado Muñoz, “Developing an orientation and cutting point determination algorithm for a trout fish processing system using machine vision,” *Computers and Electronics in Agriculture*, vol. 162, pp. 613–629, 2019.
- [124] M. Abdelhamid, Y. A. Sudnik, H. Alshinayyin, and F. Shaaban, “Chlorophyll fluorescence for classification of tomato fruits by their maturity stage,” in *Proceedings of the Paper presented at the E3S Web of Conferences*, 2020.
- [125] J. Zhuang, Y. Wang, Y. Chi et al., “Drought stress strengthens the link between chlorophyll fluorescence parameters and photosynthetic traits,” *PeerJ*, vol. 8, Article ID e10046, 2020.
- [126] F. Moradinezhad, F. Setayesh, S. Mahmoodi, and M. Khayyat, “Physicochemical properties and nutritional value of jujube (*Ziziphus jujuba* Mill.) fruit at different maturity and ripening stages,” *International Journal of Horticultural Science and Technology*, vol. 3, no. 1, pp. 43–50, 2016.
- [127] W. Pan, W. Ma, J. Zheng, and S. Geng, “Chlorophyll fluorescence of tobacco seeds as marker of seed maturity,” *Southwest China Journal of Agricultural Sciences*, vol. 29, no. 4, pp. 966–969, 2016.
- [128] H. Lou, Y. Hu, L. Zhang, P. Sun, and H. Lu, “Nondestructive evaluation of the changes of total flavonoid, total phenols, ABTS and DPPH radical scavenging activities, and sugars during mulberry (*Morus alba* L.) fruits development by chlorophyll fluorescence and RGB intensity values,” *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, vol. 47, no. 1, pp. 19–24, 2012.
- [129] M. Khojastehnazhand, V. Mohammadi, and S. Minaei, “Maturity detection and volume estimation of apricot using image processing technique,” *Scientia Horticulturae*, vol. 251, pp. 247–251, 2019.
- [130] M. Fashi, L. Naderloo, and H. Javadikia, “The relationship between the appearance of pomegranate fruit and color and size of arils based on image processing,” *Postharvest Biology and Technology*, vol. 154, pp. 52–57, 2019.
- [131] V. Mohammadi, K. Kheiralipour, and M. Ghasemi-Varnamkhasi, “Detecting maturity of persimmon fruit based on image processing technique,” *Scientia Horticulturae*, vol. 184, pp. 123–128, 2015.
- [132] A. Nasiri, A. Taheri-Garavand, and Y.-D. Zhang, “Image-based deep learning automated sorting of date fruit,” *Postharvest Biology and Technology*, vol. 153, pp. 133–141, 2019.
- [133] A. Jahanbakhshi and K. Kheiralipour, “Carrot sorting based on shape using image processing, artificial neural network, and support vector machine,” *Journal of Agricultural Machinery*, vol. 9, no. 2, pp. 295–307, 2019.
- [134] S. Jarolmasjed, C. Z. Espinoza, S. Sankaran, and L. R. Khot, “Postharvest bitter pit detection and progression evaluation in ‘Honeycrisp’ apples using computed tomography images,” *Postharvest Biology and Technology*, vol. 118, pp. 35–42, 2016.
- [135] J. I. Larada, G. J. Pojas, and L. V. V. Ferrer, “Postharvest classification of banana (*Musa acuminata*) using tier-based machine learning,” *Postharvest Biology and Technology*, vol. 145, pp. 93–100, 2018.
- [136] D. F. A. Riza, T. Suzuki, Y. Ogawa, and N. Kondo, “Diffuse reflectance characteristic of potato surface for external defects discrimination,” *Postharvest Biology and Technology*, vol. 133, pp. 12–19, 2017.
- [137] A. Beyaz, M. T. Özkaya, and D. İcen, “Identification of some Spanish olive cultivars using image processing techniques,” *Scientia Horticulturae*, vol. 225, pp. 286–292, 2017.
- [138] H. Azarmdel, A. Jahanbakhshi, S. S. Mohtasebi, and A. R. Muñoz, “Evaluation of image processing technique as an expert system in mulberry fruit grading based on ripeness level using artificial neural networks (ANNs) and support vector machine (SVM),” *Postharvest Biology and Technology*, vol. 166, Article ID 111201, 2020.
- [139] G. Somaratne, M. M. Reis, M. J. Ferrua et al., “Mapping the spatiotemporal distribution of acid and moisture in food structures during gastric juice diffusion using hyperspectral imaging,” *Journal of Agricultural and Food Chemistry*, vol. 67, no. 33, pp. 9399–9410, 2019.
- [140] Y. Ji, L. Sun, Y. Li et al., “Non-destructive classification of defective potatoes based on hyperspectral imaging and support vector machine,” *Infrared Physics & Technology*, vol. 99, pp. 71–79, 2019.
- [141] Y. Shao, Y. Wang, G. Xuan et al., “Assessment of strawberry ripeness using hyperspectral imaging,” *Analytical Letters*, vol. 54, no. 10, pp. 1–14, 2020.
- [142] J. Zabalza, J. Ren, M. Yang et al., “Novel folded-PCA for improved feature extraction and data reduction with hyperspectral imaging and SAR in remote sensing,” *ISPRS Journal of Photogrammetry and Remote Sensing*, vol. 93, pp. 112–122, 2014.
- [143] L. Yang, L. Meng, H. Gao et al., “Heavy metal detection in mulberry leaves: laser-induced breakdown spectroscopy data,” *Data in Brief*, vol. 33, Article ID 106483, 2020.
- [144] L. Huang, Y. Zhou, L. Meng, D. Wu, and Y. He, “Comparison of different CCD detectors and chemometrics for predicting total anthocyanin content and antioxidant activity of mulberry fruit using visible and near infrared hyperspectral imaging technique,” *Food Chemistry*, vol. 224, pp. 1–10, 2017.
- [145] L. Yang, H. Gao, L. Meng et al., “Nondestructive measurement of pectin polysaccharides using hyperspectral imaging in mulberry fruit,” *Food Chemistry*, vol. 334, Article ID 127614, 2021.